

-147-

Any 14 - or 15-membered-ring macrolides have not been reported to be effective against non-small cell lung cancers.

However, several chemotherapeutic agents have been shown to be efficacious against NSCLC. Preferred chemotherapeutic agents that can be used in the present invention against NSCLC include etoposide, carboplatin, methotrexate, 5-Fluorouracil, epirubicin, doxorubicin, taxol, inhibitor of normal mitotic activity; and cyclophosphamide. Even more preferred chemotherapeutic agents active against NSCLC include cisplatin, ifosfamide, mitomycin C, epirubicin, vinblastine, and vindesine.

Other agents that are under investigation for use against NSCLC include: camptothecins, a topoisomerase 1 inhibitor; navelbine (vinorelbine), a microtubule assembly inhibitor; gemcitabine, a deoxycytidine analogue; fotemustine, a nitrosourea compound; and edatrexate, a antifol.

The overall and complete response rates for NSCLC has been shown to increase with use of combination chemotherapy as compared to single-agent treatment. Haskel CM: Chest. 99: 1325, 1991; Bakowski MT: Cancer Treat Rev 10:159, 1983; Joss RA: Cancer Treat Rev 11:205, 1984.

A preferred therapy for the treatment of NSCLC is a combination of therapeutically effective amounts of one or more COX-2 inhibitors in combination with the following combinations of antineoplastic agents: 1) ifosfamide, cisplatin, etoposide; 2) cyclophosphamide, doxorubicin, cisplatin; 3) ifosfamide, carboplatin,

-148-

etoposide; 4) bleomycin, etoposide, cisplatin;  
5) isofamide, mitomycin, cisplatin; 6) cisplatin,  
vinblastine; 7) cisplatin, vindesine; 8) mitomycin C,  
vinblastine, cisplatin; 9) mitomycin C, vindesine,  
5 cisplatin; 10) isofamide, etoposide; 11) etoposide,  
cisplatin; 12) isofamide, mitomycin C; 13) flurouracil,  
cisplatin, vinblastine; 14) carboplatin, etoposide; or  
radiation therapy.

Accordingly, apart from the conventional concept of  
10 anticancer therapy, there is a strong need for the  
development of therapies practicably effective for the  
treatment of non-small cell lung cancers.

#### Small Cell Lung Cancer

Approximately 15 to 20 percent of all cases of lung  
15 cancer reported worldwide is small cell lung cancer  
(SCLC). Ihde DC: Cancer 54:2722, 1984. Currently,  
treatment of SCLC incorporates multi-modal therapy,  
including chemotherapy, radiation therapy and surgery.  
Response rates of localized or disseminated SCLC remain  
20 high to systemic chemotherapy, however, persistence of  
the primary tumor and persistence of the tumor in the  
associated lymph nodes has led to the integration of  
several therapeutic modalities in the treatment of SCLC.

A preferred therapy for the treatment of lung  
25 cancer is a combination of therapeutically effective  
amounts of one or more COX-2 inhibitors in combination  
with the following antineoplastic agents: vincristine,  
cisplatin, carboplatin, cyclophosphamide, epirubicin  
(high dose), etoposide (VP-16) I.V., etoposide (VP-16)  
30 oral, isofamide, teniposide (VM-26), and doxorubicin.  
Other preferred single-agents chemotherapeutic agents

-149-

that may be used in the present invention include BCNU (carmustine), vindesine, hexamethylmelamine (altretamine), methotrexate, nitrogen mustard, and CCNU (lomustine). Other chemotherapeutic agents under investigation that have shown activity against SCLC include irroplatin, gemcitabine, lonidamine, and taxol. Single-agent chemotherapeutic agents that have not shown activity against SCLC include mitoguazone, mitomycin C, aclarubicin, diaziquone, bisantrene, cytarabine, idarubicin, mitomxantrone, vinblastine, PCNU and esorubicin.

The poor results reported from single-agent chemotherapy has led to use of combination chemotherapy.

A preferred therapy for the treatment of NSCLC is a combination of therapeutically effective amounts of one or more COX-2 inhibitors in combination with the following combinations of antineoplastic agents: 1) etoposide (VP-16), cisplatin; 2) cyclophosphamide, adriamycin [(doxorubicin), vincristine, etoposide (VP-16)]; 3) Cyclophosphamide, adriamycin(doxorubicin), vincristine; 4) Etoposide (VP-16), ifosfamide, cisplatin; 5) etoposide (VP-16), carboplatin; 6) cisplatin, vincristine (Oncovin), doxorubicin, etoposide.

Additionally, radiation therapy in conjunction with the preferred combinations of COX-2 inhibitors and/or systemic chemotherapy is contemplated to be effective at increasing the response rate for SCLC patients. The typical dosage regimen for radiation therapy ranges from 40 to 55 Gy, in 15 to 30 fractions, 3 to 7 times week. The tissue volume to be irradiated is determined by